

The topology of events

Vipul Periwal

Department of Physics, Princeton University, Princeton, New Jersey 08544

A matrix model is constructed to compute characteristic numbers of the space of subsets of $\mathbf{R}^d$ with N elements. This matrix model is found to be a constrained null dimensional reduction to a point of a Yang-Mills theory with anticommuting matter in $d + 2$ dimensions. The constraints and equations of motion are similar to those found by Nishino and Sezgin in their $10 + 2$ dimensional supersymmetric Yang-Mills equations. It is conjectured that the $d = 10$ topological model is a twisted form of the Nishino-Sezgin model, dimensionally reduced to a point, due to $SO(8)$ triality. Topological matrix models are constructed for non-commutative spaces, entirely in terms of algebras associated with such spaces. These models exhibit degrees of freedom analogous to massive modes in string theory.

An amusing feature of much recent work (reviewed in [1–3]) on matrix models of fundamental interactions is the growing interest in noncommutative geometry, and the possibility that the spacetime continuum of common experience is replaced by something described by matrix coördinates, first observed in [4].

I consider in this letter an *a priori* different problem, that of calculating some characteristic numbers of the space of finite subsets of cardinality N in some manifold, M . I show that, when $M = \mathbf{R}^d$, this problem is solved by a topological matrix model that resembles a dimensional reduction to a point of a Yang-Mills model in $d + 2$ dimensions. I find that the features of this topological matrix model are in close correspondence to the dimensional reduction of the Nishino-Sezgin supersymmetric Yang-Mills equations [5,6], which I proposed some time ago as a fundamental description of M theory [7]. This leads to the conjecture that precisely when $d = 10$, the topological model can be twisted to give rise to the (reduced) Nishino-Sezgin model, on account of $SO(8)$ triality. Thus, consistent with the spin-statistics theorem, only when $d = 9 + 1$ does the topological model possess a sensible spacetime interpretation. If this conjecture is true, non-renormalization properties of the reduced supersymmetric Yang-Mills theory would follow for a restricted class of observables from the topological invariance of the twisted theory.

In this construction that the reduced Yang-Mills form of the action appears because the functions on $\mathbf{R}^d$ form a commutative algebra. It is then a simple matter to find a topological model appropriate for describing N points in an arbitrary space, in terms of the (possibly non-commutative) algebra associated with the space [8]. Some features of this algebraic construction are taken from Ref. [9]. There appear additional degrees of freedom, even in the case of $\mathbf{R}^d$, that may be analogous to the massive modes in string theory.

Lastly, I point out that the appropriate mathematical structure underlying these constructions seems to be a variant of operads [10]. Operads appear, of course, in conformal field theory [11] and in string field theory [12] already, though the structures that appear here seem distinct. This paper contains only a physical approach to the problem. I hope to return to the mathematical framework elsewhere.

Consider a set with N indistinguishable points, B_N . The standard algebra of functions to associate with B_N is the cross product algebra obtained from the Cartesian product of N copies of $\mathbf{R}$ by the action of the symmetry group on N letters. Another algebra one could associate with B_N , is the algebra of $N \times N$ Hermitian matrices, H_N , up to unitary equivalence—the eigenvalues of the matrices, up to permutations, are the analogues of the values of a function on the N points. This is the algebra that arises naturally in Witten’s description [4] of D-instantons. D-instantons have been studied in Ref. [13].

Now consider the space of maps from B_N to $\mathbf{R}^d$. In terms of coördinates on $\mathbf{R}^d$, each map $x^i : \{1, \dots, N\} \rightarrow \{x^i(1), \dots, x^i(N)\}$, so this space is equivalently described as the space of subsets of $\mathbf{R}^d$ with N elements. Such maps give rise to obvious homomorphisms from the continuous functions on $\mathbf{R}^d$, $\mathcal{C}(\mathbf{R}^d) \equiv \mathcal{C}(\mathbf{R})^{\otimes d}$ to H_N , by pullback: Every element $f \in \mathcal{C}(\mathbf{R})^{\otimes d}$ gets mapped to $x^*f(\{1, \dots, N\}) \equiv \text{diag}\{f(x(1)), f(x(2)), \dots, f(x(N))\}$, up to equivalence by unitary conjugation.

To start with, we will suppose that an element of $\text{Hom}(\mathcal{C}(\mathbf{R})^{\otimes d}, H_N)$ is specified by its action on the generators of $\mathcal{C}(\mathbf{R})^{\otimes d}$, which we take to be the coördinate functions, x^i . Thus, an element of $\text{Hom}(\mathcal{C}(\mathbf{R})^{\otimes d}, H_N)$ is specified by d commuting elements of H_N , X^i . We want to compute topological invariants of $\text{Hom}(\mathcal{C}(\mathbf{R})^{\otimes d}, H_N)$. To this end, we want to construct a measure to integrate over d arbitrary elements of H_N , which is concentrated on configurations of d commuting matrices. We want the measure to be invariant under unitary conjugation, in order that we describe only unordered sets.

There is a standard procedure for constructing such measures, extracted from Witten’s work [14] on a quantum field-theoretic interpretation of the Donaldson invariants. One introduces a set of d matrices with anticommuting entries, which we label ψ^j . Further we define an anticommuting operator s acting on X^i, ψ^j :

$$sX^i = \psi^i, \quad s\psi^i = i[\phi, X^i], \quad s\phi = 0 \quad (1)$$

where ϕ is a matrix with commuting entries. We introduce a set of matrices with anticommuting entries, b^{ij} (antisymmetric in i, j) and L , and a matrix with commuting entries, K , on which s acts as

$$sb^{ij} = i[X^i, X^j], \quad sK = L, \quad sL = i[\phi, K]. \quad (2)$$

For orientation, note that s is the equivariant BRS operator for a topological shift symmetry, b^{ij} will play the rôle of the antighost for the constraint of interest $[X^i, X^j] = 0$, ψ^i is the ghost for the topological shift symmetry, and K is the antighost for the constraint $[X_i, \psi^i] = 0$. The latter constraint is standard in constructions of topological field theories, and ensures that zero-modes of ψ^i are in 1–1 correspondence with moduli deformations. To see this, note that if $\{X^i\}$ is a solution of $[X^i, X^j] = 0$, $\{X^i + i[X^i, E]\}$ is also a solution, for any matrix E . Thus the ψ equations of motion will automatically have solutions of the form $i\epsilon[X^i, E]$, where ϵ is an anticommuting number. If we demand $[X_i, \psi^i] = 0$, such solutions are eliminated, leaving only the desired solutions.

Relabel the matrices as follows: Let $I, J \in \{1, \dots, d, +, -\}$, and define $b^{+I} \equiv \psi^I$, with $\psi^- \equiv L$. By antisymmetry, $b^{++} = 0$, and we constrain $b^{-j} = 0$. Further, define $X^+ \equiv \phi$, and $X^- \equiv K$. We see now that the action of s given in eq.'s 1,2 may be written more succinctly as

$$sX^I \equiv b^{+I}, \quad sb^{+I} = i[X^+, X^I], \quad sb^{ij} = i[X^i, X^j]. \quad (3)$$

While the matrices are naturally organized in this sort of way, there is clearly no $d + 2$ dimensional Lorentz-like symmetry in the action of s , since the $+$ index appears explicitly in eq. 3. With the usual ghost numbers, we see that there is a simple rule: $gh(X^i) = 0$, $gh(X^\pm) = \pm 2$, and $gh(b^{ij}) = -1$, $gh(b^{+i}) = +1$, $gh(b^{+-}) = -1$. Thus an upper $\pm$ index corresponds to an additional ghost number ± 2 . It seems natural to think of this ghost number as the analogue of Ramond-Ramond charge. It would be interesting to relate this to the work of Berkovits [15].

We will need a metric for raising and lowering indices. Let us suppose that the metric on $\mathbf{R}^d$ is Minkowskian of some signature $(p, d - p)$ —different choices are irrelevant for the topological theory, but are relevant for the physical theory (see below). We extend the inner product to include the $\pm$ indices so that $(v^i, v^+, v^-) \cdot (w^i, w^+, w^-) = v^i w_i + v^+ w^- + v^- w^+$. Now consider the gauge fermion

$$\Psi = i \text{tr} (b_{ij}[X^i, X^j] + 2X^-[X_i, b^{+i}]) = i \text{tr} (b_{ij}[X^i, X^j] + 2b^{+i}[X^-, X_i]). \quad (4)$$

We have

$$s\Psi = i \text{tr} \left(i[X_i, X_j][X^i, X^j] + 2i[X^-, X_i][X^+, X^i] + b_{ij}([X^j, b^{+i}] - [X^i, b^{+j}]) + 2b^{+-}[X_i, b^{+i}] - 2b^{+i}[X^-, b^{+i}] \right).$$

Let $\Delta S \equiv \frac{i}{2} \text{tr}(b_{ij}[X^+, b^{ij}])$. Define the complete action $S \equiv s\Psi + \Delta S$. All terms with anticommuting fields have an upper $+$ index, the analogue of a chiral projection for fermion terms. Note that $s^2 \cdot = i[X^+, \cdot]$, and computing $sS = s^2\Psi + s\Delta S$, we find $sS = 0$, using the b_{ij} equation of motion. s is therefore a scalar supersymmetry, squaring to a bosonic field-dependent infinitesimal $U(N)$ transformation, the appropriate structure for equivariant cohomology. It is easy to write S in a manifestly off-shell supersymmetric form by introducing a Lagrange multiplier, Λ^{ij} .

Notice that the terms in S quartic in X are not exactly in correspondence with a dimensional reduction of a Yang-Mills action in $d + 2$ dimensions. One mismatch is the absence of the term $2\text{tr}[X^+, X^-]^2$, and the other mismatch is the coefficient of the term $\text{tr}[X^+, X^i][X^-, X_i]$. The first mismatch could be easily remedied by adding a term $2i\text{tr}b^{+-}[X^-, X^+]$ to Ψ in eq. 4. This term has the correct ghost number -1 to be added to Ψ , and leads merely to a small modification of the $[X_i, b^{+i}] = 0$ constraint. It also does not change the b_{ij} equation of motion, and therefore is consistent with the supersymmetry. However, I do not add such a term because the action S is easier to study as it stands, since $X^\pm$ appear linearly. The remaining mismatch is due to the fact that there are no fields b^{-j} . The terms with anticommuting fields can be written as

$$\text{tr} \left(b_{[IJ} n_{K]} [X^I, b^{JK}] \right), \quad \text{with } n_J \equiv \delta_{J+}, \quad (5)$$

except for the coefficient of the term in ΔS .

The condition $s^2 = 0$ is the analogue in the present case of the BPS condition $\det\{Q_\alpha, Q_\beta\} = 0$. Thus, imposing the constraint $[X^+, X^I] = 0$, we find

$$[X_j, [X^j, X^i]] + \frac{i}{2} \{b^{+j}, b^i_j - b^{+-} \delta^i_j\} = 0, \quad [X_i, [X^i, X^-]] - \frac{i}{4} \{b^{ij}, b_{ij}\} = 0. \quad (6)$$

Let us compare this set-up to the equations and constraints found by Nishino and Sezgin [5] in their construction of supersymmetric Yang-Mills equations in $10 + 2$ dimensions. Consider a vector field X^I and a positive chirality

Majorana-Weyl fermion λ in $10 + 2$ dimensions. Dimensionally reduced to $0 + 0$ dimensions [7], their constraints [5] are

$$[X^+, X^I] = 0 = [X^+, \lambda], \quad \Gamma^+ \lambda = 0, \quad (7)$$

and their bosonic equation of motion is

$$[X_J, [X^J, X_{[K}]]n_L] + \frac{1}{4}\{\bar{\lambda}, \Gamma_{KL}\lambda\} = 0. \quad (8)$$

We see that supersymmetric solutions in the topological theory constructed above satisfy the same constraint of commuting with X^+ , and that eq. 6 has precisely the same bosonic terms as eq. 8. $\Gamma^+ \lambda = 0$ is the analogue of $b^{-j} = 0$, rather than $b^{+j} = 0$, since Γ^I changes chirality. I do not understand the exact mapping of anticommuting degrees of freedom, which depends on the signature of the metric. However, Nishino [6] has given an action underlying the Nishino-Sezgin equations [5], and this action has a fermion term $\text{tr}(\bar{\lambda}\Gamma^{IJ}[X_I, \lambda]n_J)$, which is similar to eq. 5.

This leads to the following conjecture: Precisely when $d = 9 + 1$, the topological model equations for supersymmetric solutions are a twisted form of the Nishino-Sezgin equations, with a twist related to $\text{SO}(8)$ triality. If this conjecture holds, it may imply that an appropriate subset of topological observables in the ‘physical’ theory (*i.e.* the theory with anticommuting fields in spinor representations) can be computed reliably in perturbation theory. The topological theories in other dimensions probably do not admit such twistings, and therefore are not of much physical interest.

Before going on to an analogous construction for a general algebra, I note that the weak coupling equations of motion for the topological model lead to formal solutions which take the form of singular differential forms [16] on the space of diagonal matrices that have poles when two (or more) of the N points coincide in $\mathbf{R}^d$. These singular differential forms are similar to those described in, for example, [17].

We see in the above topological construction for $\mathbf{R}^d$ hints of the general structure of supersymmetric matrix models [1–3]. It is natural to ask what the analogous construction would be for a general algebra. I discuss briefly the general structure of such spaces of algebra homomorphisms. Suppose the structure constants of an algebra A (resp. B) are $f_\alpha^{\beta\gamma}$ (resp. g_a^{bc}) with respect to a basis $\{s^\alpha\}$ (resp. $\{r^a\}$). Then a linear map $M : A \rightarrow B$ which takes $s^\alpha \mapsto S^\alpha_a r^a$ is an algebra homomorphism if

$$S^\alpha_a S^\beta_b g_c^{ab} = S^\gamma_c f_\gamma^{\alpha\beta}. \quad (9)$$

I have replaced x^i in the previous discussion with s^α in order to emphasize that this is a much more general setup, involving not just some set of generators of the algebra, but *every* element of a basis for the vector space structure underlying the algebra. We ask what model results when we use eq. 9 as the constraint equation, instead of $[X^i, X^j] = 0$. Of course, *if* M is a homomorphism, it is determined completely by its action on generators of A , so one could restrict the use of eq. 9 only to indices α, β corresponding to such generators [9]. Not only is such a restricted subset of equations difficult to describe in an invariant way, but we have constructed above a measure on *arbitrary* linear maps, which is only peaked on the set of homomorphisms.

When the algebra B is H_N , eq. 9 can be written as $S^\alpha S^\beta = S^\gamma f_\gamma^{\alpha\beta}$, with matrix multiplication in H_N automatically incorporating the rôle of g_a^{bc} in eq. 9. We introduce a BRS operator s , and ghost matrices with anticommuting entries $b^{+\alpha}$, such that

$$sS^\alpha = b^{+\alpha}, \quad b^{+\alpha} = i[X^+, S^\alpha], \quad sX^+ = 0, \quad (10)$$

in analogy with eq. 1. The number of matrices introduced here is infinite, even for $\mathbf{R}^d$. Thus, for $\mathbf{R}^d$, if we introduce a matrix X^1 for the coördinate function x^1 , we introduce an *independent* matrix for the function $(x^1)^2$. The principle is that we are integrating over *all* linear maps between the vector spaces underlying the algebras A and H_N , with the action constraining the linear maps to be algebra homomorphisms. Thus, it is logically necessary to map $(x^1)^2$ to a matrix which is *a priori* independent of $(X^1)^2$. These may be the analogues of massive modes in string theory.

The antighosts, $b^{\alpha\beta}$, matrices with anticommuting entries, are paired with Lagrange multipliers, $\Lambda^{\alpha\beta}$, which are matrices with commuting entries:

$$sb^{\alpha\beta} = \Lambda^{\alpha\beta}, \quad s\Lambda^{\alpha\beta} = i[X^+, b^{\alpha\beta}], \quad sX^+ = 0. \quad (11)$$

To proceed further, suppose that A is equipped with an inner product—for example, if A has a trace, we could use $\text{Tr}_A(s^\alpha s^\beta) \equiv h^{\alpha\beta}$ as the inverse of a metric. The observables of the topological theory will be independent of this inner product, but possible twists to get physical fermionic fields in spinor representations will depend on the choice of inner product. We use the inner product to raise and lower indices, and introduce $X^- \equiv K, b^{+-} \equiv L$, as in eq. 2.

The gauge fermion

$$\Psi_A \equiv 2\text{tr} \left[b_{\alpha\beta} \left(S^\alpha S^\beta - S^\gamma f_\gamma^{\alpha\beta} - \frac{1}{2} \Lambda^{\alpha\beta} \right) + ib^{+\alpha} [X^-, S_\alpha] \right] \quad (12)$$

leads to an action, $S_A \equiv s\Psi_A$, that after integrating over Λ , has terms quadratic, cubic and quartic in S^α , somewhat reminiscent of string field theory. It is instructive to see how the $[X^i, X^j]^2$ term arises for $\mathbf{R}^d$ by integrating out the mode corresponding to $x^i x^j \equiv x^j x^i$. Thus, the Yang-Mills form of matrix model actions for $\mathbf{R}^d$ is an artifact of the fact that we are considering an algebra $A \equiv \mathcal{C}(\mathbf{R})^{\otimes d}$ that happens to be commutative. The action S_A is the appropriate general form—it is important to note that minimal assumptions were made about the properties of A . (In particular, it may be of interest to take A to be a Poisson algebra, possibly associated with a symplectic structure on a manifold, in which case $S^\alpha S^\beta$ is replaced by $[S^\alpha, S^\beta]$ in eq. 12, with all else unchanged.) Integrating out appropriate S^α will reduce the functional integral to a more nonlinear form, expressed in terms of generators and relations for the algebra, A . Note especially that there seems to be a clear distinction between $X^\pm$ and S^α in S_A .

Even if the conjecture stated earlier can be proved, what does all this have to do with physics? First, we need to understand what restrictions the existence of a twisted spinorial interpretation for anticommuting fields places on the pairs $A, h_{\alpha\beta}$. A possible line of attack is to take the bosonic terms of S_A and work out conditions for the existence of a supersymmetric extension. It may be of interest to consider the work of Baulieu, Green and Rabinovici [18] in this context. If one could obtain such an understanding, I suggest the following interpretation: Physical observations involve a very large number of ‘events’. When we consider amplitudes for large numbers of events, we find that there are saddlepoints in the matrix integral that can be interpreted as the propagation of particles or extended objects from a spacetime perspective. Nevertheless, the underlying theory may just be a ‘theory of events’, with conditional amplitudes calculated from matrix integrals. What is interesting is that the matrix integral automatically includes matrices which naïvely correspond to two additional dimensions, which may perhaps be related to F theory compactifications [19]. It may be interesting to consider a topological matrix model for configurations of N lines (or other extended objects), as well.

I am grateful to R. von Unge for valuable comments. This work was supported in part by NSF grant PHY96-00258.

-
- [1] T. Banks, *Matrix Theory*, hep-th/9710231
 - [2] W. Taylor III, *Lectures on D-branes, gauge theory and M(atrices)*, hep-th/9801182
 - [3] A. Sen, *An introduction to non-perturbative string theory*, hep-th/9802051
 - [4] E. Witten, Nucl. Phys. B460, 335 (1995)
 - [5] H. Nishino and E. Sezgin, Phys. Lett. B388, 569 (1996)
 - [6] H. Nishino, *Lagrangian and covariant field equations for supersymmetric Yang-Mills theory in 12d*, hep-th/9710141
 - [7] V. Periwal, Phys. Rev. D55, 1711 (1997)
 - [8] A. Connes, *Noncommutative geometry*, Academic Press, 1994
 - [9] V. Periwal, *Pointless strings*, Ph.D. thesis, Princeton University (1988)
 - [10] F.R. Cohen, J. Pure App. Alg. 100, 19 (1995) (and references therein)
 - [11] T. Kimura, J. Stasheff and A.A. Voronov, in *The Gelfand Mathematics Seminars, 1993-1994*, (J. Lepowsky and M.M. Smirnov, eds.), Birkhäuser, 1996
 - [12] B. Zwiebach, Nucl. Phys. B390, 33 (1993); T. Kimura, J. Stasheff and A.A. Voronov, Comm. Math. Phys. 171, 1 (1995)
 - [13] M.B. Green and P. Vanhove, Phys. Lett. B408, 122 (1997); M.B. Green and M. Gutperle, Nucl. Phys. B498, 195 (1997); Phys. Lett. B398, 69 (1997); *D-particle bound states and the D-instanton measure*, hep-th/9711107; I. Chepelev and A.A. Tseytlin, Nucl. Phys. B511, 629 (1998); A. Kehagias and H. Partouche, *D-instanton corrections as (p,q) -string effects and non-renormalization theorems*, hep-th/9712164
 - [14] E. Witten, Comm. Math. Phys. 117, 353 (1988)
 - [15] N. Berkovits, Phys. Rev. Lett. 79, 1813 (1997)
 - [16] The anticommuting fields do *not* vanish for such solutions, so these solutions should not be confused with the supersymmetric configurations discussed above.
 - [17] V.I. Arnold, Mat. Zametki (Math. Notes) 5, 227 (1969)
 - [18] L. Baulieu, M. Green and E. Rabinovici, Phys. Lett. B386, 91 (1996)
 - [19] M. Blencowe and M. Duff, Nucl. Phys. B310 (1988) 387; C.M. Hull, Nucl. Phys. B468 (1996) 113; C. Vafa, Nucl. Phys. B469 (1996) 403; A.A. Tseytlin, Nucl. Phys. B469 (1996) 51, hep-th/9612164; D. Kutasov and E. Martinec, Nucl. Phys. B477 (1996) 652, hep-th/9612102; D. Kutasov, E. Martinec and M. O’Loughlin, Nucl. Phys. B477 (1996) 675; I. Bars, Phys. Rev. D54 (1996) 5203, hep-th/9607112, hep-th/9610074; D. Jatkar and K. Rama, Phys. Lett. B388 (1996) 45